

APPLE MAGGOT TRAP

Effective - Safe - Non-Toxic!

The SpringStar Apple Maggot trap includes our one-season, controlled-release lure system. Do not open the plastic vial - attractant is released through the plastic. Place traps in trees 1 or 2 weeks before bud break, in early spring. Hang the trap 1 to 2 feet inside the tree canopy, near apples. Check for flies every 2 to 3 days and, if necessary, apply additional appropriate treatments to control flies. Re-glue trap when full of insects, when it becomes dusty, or **at least every 4 weeks.**

APPLE MAGGOT FACTS


Apple Maggot is a type of fruit fly that attacks apples, blueberries, hawthorn, plums, pears and cherries. It causes extensive fruit damage. The fly is smaller than a housefly and appears throughout the summer. Most fruits are attacked by flies that lay their eggs during late spring to early summer. Traps should be placed in mid-April, before apple fruiting, and left out until harvest.

8 1999 SpringStar

 **SpringStar USA Inc.**
Woodinville, WA 98072
800 769-1043

APPLES
PEARS
PLUMS
CHERRIES
BLUEBERRIES

Bonus: 3 Traps
+ 3 FREE
Lures



OAK STUMP
F A R M S

EASY ASSEMBLY

Assemble red trap spheres. They only go together one way! Attach hanger on one end of trap. Squeeze glue onto trap from tube. Spread about a 1/16" thick layer. Open white packet and attach Apple Maggot Lure to hanger with a twist tie. Hang the trap in your fruit tree. In case of contact, remove excess glue with vegetable oil.



Assemble Traps



Apply Glue



Spread Glue



Attach Lure

TO MONITOR APPLE MAGGOTS use all 3 traps in an orchard.
TO CONTROL APPLE MAGGOTS use 2 traps per dwarf tree and 3 traps per large tree.

Safe for organic uses, contains no pesticides. Place out of reach of children and pets.

Made in USA,
Model No. 510

